

la connectivité sédimentaire et la connectivité des environnements proglaciaires : relation spatio-temporelle de la livraison en sédiments depuis les pentes

Sediment connectivity and connectivity of proglacial environments: Spatio-temporal pattern of sediment deliver from hillslopes coupling to proglacial margins

Mathilde BAYENS (Institut Idyst, Université de Lausanne, Suisse), **Stuart LANE** (Institut Idyst, Université de Lausanne, Suisse)

1. Sediment connectivity and connectivity of proglacial environments: Spatio-temporal pattern of sediment deliver from hillslopes coupling to proglacial margins

In a context of climate change, water yield, sediment supply and transport capacity have been increasing since the early 1980s in Alpine catchments experiencing rapid glacier recession (Micheletti and Lane, 2016; Lane et al., 2017). Over the next few decades the increasing extent of deglaciated terrain will interact with changing magnitude and frequency of sediment mobilizing extreme events (e.g. debris flows) (Rebetez et al., 1997; Ravanel and Deline, 2011; Fischer et al., 2012; Stoffel and Huggel, 2012; Hirschberg et al., 2021).

In permafrost-dominated regions rising temperatures are also changing the response of slope stability to extreme precipitation events (Rebetez et al., 1997). It is not only the frequency of the events that is likely to change but also the seasonality as climate warming reduces winter snow accumulation and leads to an earlier onset of snow-free conditions (IPCC, 2021). In the same way, even if the overall frequency of debris flows events remains low, magnitude may increase if larger volumes of sediment are mobilized (Stoffel et al., 2014; Hirschberg et al., 2021). There is emerging evidence that these changes can be seen in increasing sediment yield from mountain basins.

What is perhaps less frequently considered in periglacial environments especially is the role played by connectivity as an influence upon the ways in which the signals of changing climate propagate through river catchments. Through erosion and deposition, sediment transport leads to modification of the landscape and thus the ease with which sediment can move through the landscape, that is the degree of connectivity. For instance, glacier recession leads to sidewall debuttressing and fall in local base level leading to intense erosional processes on sidewalls (e.g. gully development) which increase potential connectivity from upstream to downstream (Lane et al., 2017; Mancini et al., 2020).

Following these processes, my PhD project focus on the relative importance of the sources, sinks and pathways of sediments in the hillslopes that form after deglaciation as compared with other drivers of sediment flux (e.g. rivers) in proglacial margins. This is being achieved by combining passive environmental seismometry to qualify (and quantify) sediment transport events, and static and dynamic connectivity analyses (Cavalli et al., 2013) to understand where sources are and the role played by connectivity in modifying source-to-sink sediment transfers.

REFERENCES

- [1] Cavalli, M., Trevisani, S., Comiti, F., & Marchi, L. (2013). Geomorphometric assessment of spatial sediment connectivity in small Alpine catchments. *Geomorphology*, 188, 31–41. <https://doi.org/10.1016/j.geomorph.2012.05.007>
- [2] Fischer, L., Purves, R. S., Huggel, C., Noetzli, J., & Haeblerli, W. (2012). On the influence of topographic, geological and cryospheric factors on rock avalanches and rockfalls in high-mountain areas. *Natural Hazards and Earth System Sciences*, 12(1), 241–254. <https://doi.org/10.5194/nhess-12-241-2012>
- [3] Hirschberg, J., Fatichi, S., Bennett, G. L., McArde, B. W., Peleg, N., Lane, S. N., Schlunegger, F., & Molnar, P. (2021). Climate Change Impacts on Sediment Yield and Debris-Flow Activity in an Alpine Catchment. *Journal of Geophysical Research: Earth Surface*, 126(1). <https://doi.org/10.1029/2020JF005739>
- [4] IPCC. (2021). 'Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (Eds.)].' Cambridge University Press. In Press.
- [5] Mancini, D., & Lane, S. N. (2020). Changes in sediment connectivity following glacial debuttressing in an Alpine valley system. *Geomorphology*, 352, 106987. <https://doi.org/10.1016/j.geomorph.2019.106987>
- [6] Micheletti, N., & Lane, S. N. (2016). Water yield and sediment export in small, partially glaciated Alpine watersheds in a warming climate: CLIMATE CHANGE CONTROL IN SMALL ALPINE WATERSHEDS. *Water Resources Research*, 52(6), 4924–4943. <https://doi.org/10.1002/2016WR018774>
- [7] Lane, S. N., Bakker, M., Gabbud, C., Micheletti, N., & Saugy, J.-N. (2017). Sediment export, transient landscape response and catchment-scale connectivity following rapid climate warming and Alpine glacier recession. *Geomorphology*, 277, 210–227. <https://doi.org/10.1016/j.geomorph.2016.02.015>
- [8] Ravanel, L., & Deline, P. (2011). Climate influence on rockfalls in high-Alpine steep rockwalls: The north side of the Aiguilles de Chamonix (Mont Blanc massif) since the end of the 'Little Ice Age'. *The Holocene*, 21(2), 357–365. <https://doi.org/10.1177/0959683610374887>
- [9] Rebetez, M., Lugon, R., & Baeriswyl, P.-A. (1997). Climatic Change and Debris Flows in High Mountain Regions: The Case Study of the Ritigraben Torrent (Swiss Alps). In H. F. Diaz, M. Beniston, & R. S. Bradley (Eds.), *Climatic Change at High Elevation Sites* (pp. 139–157). Springer Netherlands. https://doi.org/10.1007/978-94-015-8905-5_8
- [10] Stoffel, M., & Huggel, C. (2012). Effects of climate change on mass movements in mountain environments. *Progress in Physical Geography: Earth and Environment*, 36(3), 421–439. <https://doi.org/10.1177/0309133312441010>
- [11] Stoffel, M., Mendlik, T., Schneuwly-Bollschweiler, M., & Gobiet, A. (2014). Possible impacts of climate change on debris-flow activity in the Swiss Alps. *Climatic Change*, 122(1–2), 141–155. <https://doi.org/10.1007/s10584-013-0993-z>